# Morphometric features of Azikheli buffalo in Swat Khyber Pakhtunkhwa, Pakistan

## Khan M<sup>1</sup>, Saleem M<sup>1</sup>, Farman Ullah<sup>2\*</sup>, Abass KS<sup>3</sup>, Khan SU<sup>4</sup>, Ahmad S<sup>5</sup>, khan S<sup>6</sup> and Salim M<sup>7</sup>

<sup>1</sup>Directorate General (Extension) Livestock and Dairy Development Department, Pakistan

<sup>2</sup>Faculty of Veterinary and Animal Sciences, National Center for Livestock Breeding Genetics and Genomics, LUAWMS, Pakistan
<sup>3</sup>Department of Physiology, Biochemistry and Pharmacology, College of Veterinary Medicine, University of Kirkuk; Kirkuk, Iraq
<sup>4</sup>Department of Veterinary Internal Medicine, Faculty of Veterinary Medicine, Gadjah Mada, University, Indonesia
<sup>5</sup>Institute of biotechnology and microbiology, Bacha Khan University, Charsadda, Pakistan

<sup>6</sup>Faculty of Veterinary and Animal Sciences, Lasbela University of Agriculture, Water and marine sciences (LUAWMS), Pakistan <sup>7</sup>Forestry and Wildlife Management Department, University of Haripur, Pakistan

#### Corresponding author: farman\_aup@yahoo.com

Received on: 19/12/2024	Accepted on: 20/04/2025	Published on: 25/04/2024

# ABSTRACT

**Aim:** Purpose of the study was to determine morphological and morphometric characteristics of Azikheli buffalo. **Method and materials:** The study was carried out on physical and morphometric characteristics, productive and reproductive performance of Azikheli buffalos and bulls in Khwazakhela, District Swat, Khyber Pakhtunkhwa, Pakistan. Morphometric measurements included heart girth, body length, height at wither, height at hipbone, head region (face length, ear length and width), horn, neck, back, rump, legs and tail.

**Results:** Azikheli buffalos were observed with significantly higher heart girth size, longer horns, longer neck, and wider face at the level of eyes than bulls, on the other hand Azikheli buffalo was observed with significantly longer bodies, longer ears, thick horns, thick neck and large hooves than buffalo. Tail of Azikheli buffalo and of bulls was above hock as cutting of switch was routine practice. Horns were flat laterally, directed backward and slightly upwards without twisting giving a sickle or semi-sickle appearance.

**Conclusion:** It was concluded that Azikheli buffalo is kept in its home tract by different social groups such as landowners, Gujars and tenants, under the different topographic conditions such as hill slopes, undulating areas and valley bottoms. Azikheli buffalo had large body size (heart girth) than bulls whereas bulls had longer body (body length) than Azikheli buffalo.

Keywords: Azikheli, buffalos breed, morphometric, Swat

**Cite This Article as**: Momen Khan, Saleem M, Farman Ullah, Abass KS, Khan SU, Ahmad S, khan S and Salim M (2025). Morphometric features of Azikheli buffalo in Swat Khyber Pakhtunkhwa, Pakistan. J. Vet. Res. Adv., 07(01): 28-39.

## Introduction

Pakistan is endowed with rich livestock genetic resources, well adapted to the local conditions. There are 15 breeds of cattle, 5 breeds of buffalo, 33 breeds of sheep and 36 breeds of goat (Government of Pakistan, 2006; Khan *et al.*, 2007a; Khan *et al.*, 2007b; Khan *et al.*,2008a: Khan *et al.*, 2008b; Ali *et al.*, 2009; Babar *et al.*, 2009; Khan and Niamatullah, 2010). There are 177.247 million buffalo head worldwide in 50 countries, of which 171 million (97 %) are found in Asia, while 5.38 million (3%) are found in rest of world. Pakistan with 29.9 million population (14 %) is the second highest buffalo inhabiting country in the world after India which has the population of 98.7 million (56 %) of the total world buffalo (Chanthalakhana and Falvey, 1999; Sethi, 2009; Government of Pakistan, 2009; Khan and Niamatullah, 2010). Buffalo is the second largest (75 million tons) source of milk supply in the world (FAO, 2004).

Interestingly Pakistan is the second largest dairy buffalo country in the world (Khan and Niamatullah, 2010) and possesses the highest buffalo milk genetic potential in the world and they contribute 70% of the total milk produced in the country (Khan, 2009) and with annual increase of buffalo population at the rate of 3 percent, buffalo population in Pakistan is 29.9 million head and being the major source of national milk yield (62 %) is an enormous dairy genetic resource of Pakistan (Government of Pakistan, 2009). Similarly, Pakistan is also second in Asia in terms of meat produced by buffalo which is evident from

**Copyright:** Khan *et al.* Open Access. This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (http://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

the fact that number of slaughtering buffalo increased from 2.18 million in 1996 to 3.34 million in 2006 showing a 53.2 percent increase during the specified period (Government of Pakistan, 2007). Buffalo breeds in Pakistan are Nili-Ravi and Kundi which are the finest dairy buffalo in the world (Cockrill, 1974).

Indigenous breeds that have been evolved and adapted from the time immemorial and exist with their own genetic makeup are disappearing by dilution and replacement (FAO/UNEP, 1992) because of new market demand, use of new breeding technology (FAO, 1992) changes in the socio-economic environments of a region (Dorji et al., 2009) and modern production techniques (FAO, 2010). Indigenous breeds were considered as inferior to the exotic and crossbreed animals but with passage of time the performance of indigenous breed reported was equal to or even better than that of exotic, improved or crossbred 2001). animals (Kohler-Rollefson, In the production system of harsh environmental conditions, the performance of local indigenous breeds is well than exotic breed in term of productivity (Anderson, 2003; Ayalew et al., 2003), although the output is low, but the inputs required is also low hence provide better financial returns to the farmer (Scarpa et al., 2003). Indigenous breeds are mainly kept in low-inputlow-output production system (Scherf et al., 2005), are hardy, disease resistant, survive on little water, scanty and poor vegetation, have tasty meat and good adaptability to various environments where modern imported exotic breeds unable to exist (Dong Xuan, et al., 2006; Kohler-Rollefson, 2009; FAO, 2010).

Local indigenous breeds those adapted to harsh environment of developing countries have not yet been sufficiently characterized and in the case of their extinction the value lost to humankind is not known (Scherf et al., 2005). It is therefore necessary first to evaluate local breeds for phenotype, special characteristics, performance, and performance potential and crossbred suitability especially in their home tracts and under existing management condition (Swaminathan, 1988; Zarate, 1996). Traditionally external or internal phenotypic characters have been adopted to ascribe a given animal to a breed (Bradley et al., 1993). Phenotypic and genetic characterization of populations, breeds and species is essential for the development of appropriate breeding strategies, sustainable use of genetic diversity, genetic conservation and assessment of genetic variability (Loftus et al., 1994; Hassen et al., 2007) and thus the variations at molecular level are based on the phenotypic variations among breeds (Babar et al., 2009). Phenotypic as well as adaptive characteristics are important in identifying breed attributes for immediate use by farming communities (Zulu, commonly 2008). The used phenotypic characteristics are morphological (physical and morphometric), productive, reproductive performance, birth and adult body weight of the animals. The breed of azikheli buffalo at different ages of various natural positions and locations of khwazakhela swat was noticed (Fig 1), while the map of the study area of azikheli Buffalo breed home tract and the encircled regions represent the sampling areas of research area (Fig 2). The labelled diagram of Azikheli buffalo showing various body parts measured in the study (Fig 3). Physical and morphological characteristics:

Ranking animal breeds in a population according to their levels of phylogenetic distribution is done because of morphological characters (Gatesy and Arctander, 2000; Ndumu et al., 2008; Duguma et al., 2010) and morphological characters evaluate breeding goals (Zechner et al., 2001; Zechner et al., 2002; Curik et al., 2003; Pretorius et al., 2004; Dario et al., 2006). In traditional system in which breeding practices are not documented, it determines such goals retrospectively (Rege, 2001), indicating animal size and weight in a simple and less expensive way (Goe et al., 2001). Type and function of animals for beef or dairy purposes and their values as potential breeding stock are also by morphological assessed characters (Brotherstone and Hill, 1991; Fernandez et al., 1997; Luo et al., 1997; Alderson, 1999; Zechner et al., 2001).

# *Morphometric characteristics*

Morphometric measurements have been used to evaluate the characteristics of the animal that may vary due to the influence of breed evolution, environment, nutrition, sex, age and physiological status and rearing system (Dia Palo *et al.*, 2001; Campanile *et al.*, 2003; Riva *et al.*, 2004; Lazzaroni and Bigini, 2005). Morphometric measurements can be easily measured under field condition and are biologically related to cost traits (Gallo *et al.*, 2001; CGRFA, 2007). Morphometric measurements have been suggested as more objective measures of

body conformation of animal (Islam et al., 1991; Janssens and Vandepitte, 2004) which is an important component of breeding and selection decision both in dairy (Schneider et al., 2003) and beef animals (Doren et al., 1989; Arango et al., 2002) and could serve as a guideline in selection of high yielding females particularly in areas where performance records are not available (Jogi and Patel, 1990). The water buffalo of Asia has been classified on morphological and behavioral criteria into two types of the River and Swamp buffalo (Lau et al., 1998). However, studies on buffalo with respect to body conformation, productive and reproductive potentialities are very limited, particularly in most local breeds which have been described as non-descript (Khan et al, 2007a: Patro et al., 2008).

In Pakistan, mostly studied buffalo breeds are Nili-Ravi and Kundi and there is 37% of the buffalo population (10.13 million) which is nondescript (Khan *et al.*, 2007b). Although these buffalo breeds have been considered nondescript, they are highly adapted to the environmental condition of the area and have great potential. Azikheli is a buffalo breed, in Swat, acclimatized to the local conditions and is reared by farming communities.

Only introductory information about the breed is available, emphasizing the scientific characterization of the breed (Jabbar, 1987). It is an important indigenous animal genetic resource of the area and got its name from its original home tract called Azikheil, one of the several tributary valleys of river Swat. The broader home tract includes the watershed of River Swat (District Swat) and River Panjkora (District Lower and Upper Dir), District Shangla, Bunair and Malakand agency. Pockets of the breed can also be found in District Mardan, Charsadda, Nowshera and Sawabi because of transhumant migration during winter season from upland pastures of District Swat and Dir. This breed needs characterization, description and improvement for sustainable future use in the adapted area. The present investigation was thus designed to study Azikheli buffalo breed with the following objective to determine morphological and morphometric characteristics of Azikheli buffalo.

# Materials and Methods

Khwazakhela valley (Azikheil) of Swat District was selected as the study area which is a central

location in the original home tract of the Azikheli buffalo (Fig 2). This is one of the most fertile valleys of Swat and lyingabout 20 kilometers from Mingora, the head quarter of Swat district. The Valley is almost 30 kilometers long and 20 kilometers wide. Major crops are wheat, rice and maize whereas; apple, fersimen (Dyospirus kaki) and shaftalo (Prunus persicum) are the main orchards. Different ethnic groups mostly occupied different ecological niches/units hence landowners were dominant in valley bottom, tenants on hill slopes and hilltops with cropping potentials and Gujar (settled livestock herders with cattle and buffalo herders; Ur-Rahim and Viaro, 2002) at hill slope and hill tops with grazing potentials. This stratification, however, was not strict/ watertight compartment and tenants and Gujars may be present in specific location in valley bottom and landowners at hill slope and hilltops. For this study the season was classified as autumn (September-October, temperature 24-26 C<sup>0</sup>), winter: (November-February, temperature zero to minus eight C<sup>o</sup>) spring (March-April, temperature 22-24 C<sup>o</sup>), summer (May-August, temperature 21-38 Co; Urdu Tourists Guide).

Morphological characteristics: Data on physical and morphometric characteristics was collected as per standard procedure of random sampling (FAO; 1986) from both Azikheli buffalo cows and buffalo bulls. i. Physical characteristics: Physical characteristics like color of the coat; forehead, eyelashes, eyes, horn, muzzle, forelegs, hind legs and hooves on each animal were recorded. ii. Morphometric characteristics: The following morphometric measurements (cm) were taken using measuring tape with animals standing on flat surface in normal position. In case of buffalo cows, measurements were taken within two to three months after parturition and adult bulls were also measured. The morphometric studies conducted were as follows which were also recorded (Fig 3).

1. Heart girth, body length, height at wither and height at hipbone.

2. Head region: Measurements taken in head region were the width of head between horns, width of head between eyes, ear length and width and face length.

3. Horns: Measurements taken were length of horn at greater and small curvatures, circumference of horn base, mid and below tip.

4. Neck: Measurements taken were neck length and circumference of neck at middle region.

#### Open access



Fig 1: The azikheli buffalo breed of different ages at various natural positions& locations of khwazakhela swat



Fig 2. Map of the study area of azikheli Buffalo home tract, the encircled regions represent the sampling areas.



Fig 3: Labelled diagram of Azikheli buffalo showing various body parts measured in this study (a) 1) width of head between horns 2) width of head between eyes 3) muzzle (b) 4) hoof circumference 5) pastern 6) rump length 7) hip height 8) loin length 9) chine 10) heart girth 11) height at wither 12) nick length 13) ear width 14) ear length 15) length of horn at greater curvature 16) horn base circumference 17) horn mid circumference 18) horn length at smaller curvature 19) horn circumference below tip 20) face length 21) nick circumference 22) height of fore leg below knee 23) height of hind leg below hock (24) body length.

5. Back: Measurements taken were loin length (length from last rib to hipbone) and chine (length from wither to the last rib).

6. Rump: Measurements taken were rump length (length from hipbone to pin bone) and width (distance between hipbones).

7. Legs: Measurements taken were height of the leg bellow knee and hock, height of pastern, hoof circumference and tail length.

*Statistical analysis:* Mean, standard error and Student's T-test were calculated for various comparisons. Chi-square test and analysis of variance was also performed using GraphPad Prism-5 (GraphPad Software, San Deigo, CA).

# **Results and Discussion**

This study was carried out on 618 Azikheli buffalo and bulls. The recorded parameters were morphological characteristics, milk production and reproductive performance. Morphometric characteristics of 135 Azikheli buffalo and bulls were studied.

Morphological characteristics: Morphological characteristics included physical and morphometric characteristics. This study was carried out on 135 buffalo of which 108 were buffalo and 27 were buffalo bull from Khwazakhela valley (Azikheil) of District Swat, Khyber Pakhtunkhwa, Pakistan.

*Morphometric characteristics:* This study was conducted on 135 buffalo out of which 108 were buffalo and 27 were buffalo bulls. Morphometric measurements taken were heart girth, body length, height at wither, height at hipbone, head, horn, neck, back, rump, legs, and tail.

Heart girth, body length, height at withers (from hoof to shoulder girdle) and height at hipbone: Mean heart girths, body length, height at withers and height at hip bone of Azikheli buffalo and bulls were recorded (Table 1). Azikheli buffalo have significantly (t (133) =4.36; P0.05). On the other hand, Azikheli buffalo bulls have significantly (t (133) = 3.28; P< 0.001) long body than buffalo. Azikheli buffalo bulls were also taller at hip bone than buffalo, but the difference was statistically not significant (t (133) = 0.75; P > 0.05).

*Head Region:* Measurements of head region like width of head between horns, width of head between eyes, ear length and width, and face length were presented (Table 2). Azikheli buffalo has a wider head region between horns and between eyes than bulls. However, the difference for the former was statistically not significant, whereas for the latter it was significant (t (133) = 3.28; P<0.001). Compared to buffalo, bulls have longer and wider ears and longerfaces than buffalo, but the difference for these measurements was also statistically not significant.

*Horns:* Mean length of the horn along the grater and smaller curvature and circumference at base, mid region and below the tip of Azikheli buffalo and buffalo bull was recorded (Table 3). Azikheli buffalo have significantly longer horns (both along greater curvature; t (133) = 3.13; P<0.001) Than Azikheli bulls. However, Azikheli bulls have significantly (t (133) = 5.13; P<0.001) thicker horns at base than Azikheli buffalo. Horn circumference at mid region and tip was not significantly different in both sexes.

*Neck Region:* Neck length and circumference of Azikheli buffalo and bulls was recorded (Table 4). Azikheli buffalo have significantly (t (133) = 2.57; P<0.05) longer but narrow neck (t (133) = 5.09; P<0.001) than Azikheli buffalo bulls.

*Back Region:* Measurement of the back region like chine length and loin length were noticed (Table 5). Azikheli buffalo bulls have longer chine than buffalo, whereas loin was longer in buffalo compared to bulls. However, there was no significant difference in chine and loin length between both sexes.

*Rump Region:* Mean rump length and rump width of Azikheli buffalo and bulls was presented (Table 6). Azikheli buffalo have shorter but wider rump than bulls but the difference for both measurements was statistically non-significant between buffalo and bulls.

*Legs Region:* Height below knee, height below hock, height of pastern and hoof circumference of Azikheli buffalo and bulls was recorded (Table 7). Azikheli bulls had longer front leg bellow knee joint and hind leg bellow hock joint compared to buffalo, but the difference was statistically not significant for both measurements. Pastern height was more in buffalo than bulls but the gender difference for the measurement was not significant. On the other hand, Azikheli bulls have significantly (t (133) = 2.5; P0.05).

Statistical analyses were applied to analyze the data. Different tests of significance were applied to ascertain significant or non-significant differences among variables under study. Where needed Chi-square test of significance, t-test, analysis of variance and regression analysis of variance were applied. Morphological features were studied

Table 1: Measurements (cm) of heart girth, body length, height at withers and height at hip bone of Azikheli buffalo and bulls in Khwazakhela valley of District Swat, KP, Pakistan.

Sex	Variables	Mean	Range
Buffalo (108)	Heart girth	191.36±1.26	149.86-223.52
	Body length	140.39±0.94	114.30-167.64
	Height at wither	131.35±0.57	119.38-147.32
	Height at hip bone	123.41±0.41	111.76-137.16
Bulls (27)	Heart girth	177.68±3.76 <sup>a</sup> * * *	136.25-205.74
	Body length	147.89±2.60 a * * *	117.58-164.25
	Height at wither	130.01±0.78	118.11-134.69
	Height at hip bone	124.08±0.67	116.84-129.56
	1 (( 1 1 1) D (0 001++++		•

Mean±SE a = Azilkheli buffalo vs bulls P<0.001\*\*\*

Table 2: Measurements (cm) of head region of Azikheli buffalo and bulls in Khwazakhela valley of District Swat, Khyber Pakhtunkhwa, Pakistan

Sex	Variables	Mean	Range
Buffalo (108)	Width of head between horns	22.74±0.27	17.78-30.48
	Width of head between eyes	20.37±0.13	17.15-22.86
	Ear length	21.39±0.18	17.78-30.48
	Ear width	16.46±0.12	15.24-20.32
	Face length	52.45±0.2	45.72-63.50
Bulls (27)	Width of head between horns	22.15±0.47	20.16-30.48
	Width of head between eyes	19.43±0.32 a * * *	17.18-23.54
	Ear length	22.80±0.23	20.32-24.86
	Ear width	16.86±0.21	15.24-19.05
	Face length	52.67±0.55	47.42-58.42
Mean±SE	a = Azilkheli buffalo vs bulls	P<0.001***	

Table 3: Measurements (cm) of horns of Azikheli buffalo and bulls in Khwazakhela valley of District Swat, Khyber Pakhtunkhwa, Pakistan

Sex	Variables	Mean	Range
Buffalo (108)	Horn greater curvature	42.52±1.02	24.13-68.58
	Horn smaller curvature	27.55±0.82	15.24-45.72
	Horn base circumference	22.60±0.22	17.78-27.94
	Horn mid region circumference	20.13±0.22	13.97-25.40
	Horn below tip circumference	7.63±0.13	05.08-12.70
Bulls (27)	Horn greater curvature	35.7±1.36 a***	24.13-50.80
	Horn smaller curvature	21.62±0.68 a * * *	13.97-27.94
	Horn base circumference	25.12±0.41 a * * *	21.45-30.48
	Horn mid region circumference	20.83±0.31	16.51-23.46
	Horn below tip circumference	7.13±0.21	05.08-08.89
Mean±SE	a = Azilkheli vs bull P<	0.001***	

Mean±SE

Table 4: Measurements (cm) of neck region of Azikheli buffalo and bulls in Khwazakhela valley of District Swat, Khyber Pakhtunkhwa, Pakistan.

Sex	Variables	Mean	Range
Buffalo (108)	Neck length	42.57±0.37	35.36-53.34
	Neck circumference	89.60±0.61	73.66-101.6
Bulls (27)	Neck length	40.50±0.57 <sup>a</sup> *	34.58-45.72
	Neck circumference	97.32±1.81 a * * *	71.12-114.3
Mean±SE a	a = Azilkheli vs bull P<0.05*	P<0.001***	

Table 5: Measurements (cm) of back region of Azikheli buffalo and bulls in Khwazakhela valley of District Swat, Khyber Pakhtunkhwa, Pakistan.

Sex	Variables	Mean	Range
Buffalo (108)	Loin length	35.97±0.36	25.40-43.18
	Chine	45.95±0.54	35.56-66.04
Bulls (27)	Loin length	34.55±0.55	28.45-43.18
	Chine	46.76±0.80	38.10-55.88

Mean±SE

P<0.001

Table 6: Measurements (cm) of rump of Azikheli buffalo and bulls in Khwazakhela valley of District Swat, Khyber Pakhtunkhwa, Pakistan

Sex	Variables	Mean	Range
Buffalo (108)	Rump length	41.45±0.31	33.02-48.26
	Rump width	51.03±0.37	41.91-58.42
Bulls (27)	Rump length	42.03±0.39	38.10-45.72
	Rump width	50.43±0.59	44.45-55.88

Mean+SE

Table 7: Measurements (cm) of Legs in Azikheli buffalo and bulls in Khwazakhela valley of District Swat, Khyber Pakhtunkhwa, Pakistan

Sex	Variables	Mean	Range
Buffalo (108)	Height of the leg below knee	30.34±0.25	25.40-41.91
	Height of the leg below hock	46.27±0.32	40.64-53.34
	Height of the pastern	5.92±0.11	02.54-70.62
	Hoof circumference	51.26±0.38	40.64-58.42
	Tail length	71.39±1.04	55.88-101.6
Bulls (27)	Height of the leg below knee	31.34±0.43	27.25-35.56
	Height of the leg below hock	47.51±0.38	43.18-50.80
	Height of the pastern	5.65±0.14	05.04-07.62
	Hoof circumference	53.53±1.00 ª *	45.72-76.20
	Tail length	67.38±1.58	53.34-76.46
Mean+SE	a = Azilkheli buffalo vs bull P-	<0.05*	

because of the reason that they constitute information regarding the identification of this breed. Unless we identify any breed the description of it shall be of no importance. Since Azikheli buffalo is a non-descript breed it was essential to describe the details of its morphological traits. Gender differences have also been analyzed. The analysis showed that Azikheli buffalo have significantly higher heart girth, longer horns, longer neck and wider face than bulls. But the bulls compared to buffalo have longer bodies, longer ears, thick horns, thick neck and large hoofs.

Azikheli buffalo is an indigenous breed of District Khyber Pakhtunkhwa. Very meager Swat, information is available on phenotypic as well productive and reproductive characteristics of this breed. The present study was conducted to investigate the physical and morphometric characteristics, and performance of Azikheli buffalo in District Swat, Khyber Pakhtunkhwa whereas morphometric measurements of different body parts were taken.

Morphometric measurements are used to evaluate the characteristics of the animal, and they vary due to the influence of breed evolution, environment, nutrition, sex, age, physiological status, rearing system and related to cost traits (Dia Palo, 2001; Campanil et al., 2003; Riva et al., 2004; Lazzaroni and Bigini, 2005; CGRFA, 2007). Morphometric measurement is the measurement of body conformation of animal (Janssens and

Vandepitte, 2004; Janseens et al., 2004) which is an important component of breeding and selection decision (Schneider et al., 2003) and could serve as a guideline in selection of high yielding females particularly in areas where performance records are not available (Jogi and Patel, 1990). Azikheli buffalo have significantly higher heart girth size, longer horns, longer neck and wider face at the level of eyes than bulls (Table 1-4). On the other hand, Azikheli bulls, compared to Azikheli buffalo, have significantly longer bodies, longer ears, thick horns, thick neck and large hooves. The body length and height at hipbone of Azikheli bulls were higher than buffalo whereas, height at wither was higher in buffalo than bulls. Soysal et al (2007) also reported higher values for body length and height at hipbone in Anatolian bull. Tail in Azikheli buffalo is above hock as cutting point of switch, is routine practice. In Nili-Ravi buffalo tail is long extending bellow hock and had a white switch compared to that in Azikheli buffalo (Vij and Tantia, 2005). Azikheli buffalo had smaller heart girth size (191.36±1.26 cm) than Nili-Ravi buffalo (215-225 cm; Khan et al., 1982; Ranjhan and Pathak, 1993; Moioli and Borghese, 2005) whereas height at wither of Azikheli buffalo (131.35±0.57cm) falls within the range of Nili-Ravi buffalo (125-135 cm Khan et al., 1982; Ranjhan and Pathak, 1993; Moioli and Borghese, 2005). However, Nili-Ravi buffalo are longer (145-149 cm; Khan et al., 1982; Ranjhan and Pathak, 1993; Moioli and Borghese, 2005) than

Azikheli buffalo (140.39±0.94). On the other hand, Azikheli bull has smaller heart girth, smaller at wither and have shorter body than Nili-Ravi bulls (225-226 cm; 135-137 cm; 159-165 cm; Khan et al., 1982; Ranjhan and Pathak, 1993; Moioli and Borghese, 2005). There is low heart girth in Azikheli buffalo compared to that in kundi buffalo (205 cm; Khan et al., 1982; Moioli and Borghese, 2005) but are taller at wither height (112-125 cm; Khan et al., 1982; Moioli and Borghese, 2005) and have longer body (137 cm; Khan et al., 1982; Moioli and Borghese, 2005) than kundi buffalo. Bulls of Azikheli buffalo breed have smaller heart girth size than Kundi bulls (217 cm; Khan et al., 1982; Moioli and Borghese, 2005). However, the height at wither and body length of Azikheli buffalo approximate that of the Kundi bull (Khan et al., 1982; Moioli and Borghese, 2005). Pandharpuri buffalo have comparable heart girth size (192.79 cm; Patil et al., 1998).and height at wither (132.92 cm; Patil et al., 1998) to that of Azikheli but the latter is longer than the former (130.02 cm; Patil et al., 1998). Gaddi buffalo have also comparable values to that of Azikheli buffalo with respect to heart girth, height at wither and body length (194.98±1.8, 131.3±1.1 cm; 141.2±1.63 cm; Kumar and Raj, 2007). Nagpuri buffalo had smaller heart girth, height at wither and body length than that of Azikheli buffalo (172.61±4.12 cm; 121.41±2.34 cm; 128.00±3.70 cm; Shrikhande et al., 1996). There was no significant difference in face length between Azikheli buffalo and bull. Ranjhan and Pathak (1993) also reported no difference in face length in Nili-Ravi buffalo (58 cm) and bull (58 cm). Rump length and rump width in both sexes were not significantly different in Azikheli breed. Ranjhan and Pathak (1993) also reported the same values for mean rump length (45 cm each for buffalo and bull) and rump width without marked variation (61.2 cm for buffalo and 62 cm for bull) in both sexes of Murrah buffalo. However Terzano et al (2007) reported that in Romania buffalo heifers rump length was significantly different in intensive feeding (34.5% mais silage, 31% hay, 13.8% maize meal, 10.9% Soya been and 9.8% wheat flour on DM basis (14% crude protein and 0.88 MFU/kg DM) as compared to pasture system (45±12% grass, 12±9% legume, 29±11% composite and 14±15% other species (0.49±0.11 MFU/kg DM, 14.6±6.0% crude protein and 22.8±3.4% crude fiber).Similarly, there was no significant difference in the height of forelegs below knees, height of the hind legs below hock and height of pastern in both sexes. Ranjhan

and Pathak (1993) also reported mean height of legs below knee in Murrah buffalo was 21.6 cm and it was 24.6 cm for Murrah bull without mentioning any statistical analysis. Based on horns, buffaloes are also divided into two groups. One group consists of the horns which are closed and set close to head and are down swept: e.g Murrah, Nili-Ravi, Mehsana, Jaffarabadi and Sambalpur and the other group consists of the horns which are sickle shaped and un-swept: e.g Bhadawari, Kalahandi, Kanara, Nagpuri, Tarai and Toda (Singh and Barwal; 2010). The Azikheli buffalo breed falls in the second group of horn shapes. Horns are flat laterally, directed backward and slightly upwards without twisting. Upwards turning is variable and gives a sickle or semi-sickle appearance to the horn in the breed under study. It is like Surti breed (Moioli and Borghese, 2005). It is different from that of Nili-Ravi and Kundi in which the horns are short and twisted/curled (Moioli and Borghese, 2005). Size of the horn of Azikheli buffalo observed in this study is smaller than Nagpuri (50-65 cm; Shrikhande et al., 1996; Moioli and Borghese, 2005), Gauli (51.82±3.28 cm; Kolt and Sadekar, 1996) and Chilika (49±0.2 cm, Patro et al., 2008), but longer than Nili-Ravi (Sukla et al., 2006) and Kundi (Moioli and Borghese, 2005). Horns of Azikheli buffalo were significantly longer than bull whereas, bulls have significantly thicker horn base circumference than buffalo. Soyal et al (2007) reported longer and thicker horn in female Anatolian buffalo. However, Roth (2004) in Asian buffalo reported smaller horn in female buffalo as compared to that of bull. This study reported information on morphometry, like leg color patterns, makes different buffalo breeds distinct from each other, perhaps this may also be used as a good identification mark.

## Conclusion

It was concluded morphological characteristics of the Azikheli buffalo in khwazakhela swat Khyber Pakhtunkhwa Pakistan. Azikheli buffalo is kept in its home tract by different social groups such as landowners, Gujars, and tenants, under the different topographic conditions such as hill slopes, undulating areas, and valley bottoms. They are currently caring more for the breed as they are primarily defendant for their livelihood on the sale of livestock and livestock products. Azikheli buffalo had large body size (heart girth) than bulls whereas bulls had longer body (body length) than Azikheli buffalo.

# Reference

- Alderson GLH (1999). The development of a system of linear measurements to provide an assessment of the type and function of beef cattle. Animal Genetic Resources Information 25: 45-56.
- Ali S, Babar ME, Akhtar P, Ahmad A, Rahman M, Sultan JI and Shaheen HL (2009). Genetic divergence and relationship among five buffalo breeds of Pakistan using a random amplified polymorphic DNA (RAPD) analysis. Pak J Zool (Suppl.9): 303- 308.
- Arango JA and Vleck LDV (2002). Size of beef Cows: early ideas, new developments. Genet and Mol Res. 1(1): 51-63.
- Ayalew W, King J, Bruns E and Rischkowsky B (2003). Economic evaluation of smallholder subsistence livestock production: lessons from an Ethiopian goat development program. Ecological Economics 45: 473-485.
- Babar MI, Husssain T, Nadeem A, Jabeen R and Javed M (2009). Genetic Characterization of Azikheli Buffalo Breed of Pakistan Using Micro-satellites DNA Markers. Pa. J Zool. (Suppl. 9): 361-366.
- Bradley DG, Maschugh DE, Meghen C, Loftus RT, Sharp DM and Cunningham EP (1993). Genetic Characterization of indigenous cattle breeds: first results and implications for genetic improvement. FAO Anim Prod Health Paper. 110:37-44.
- Brotherstone S and Hill WG (1991). Dairy herd life in relation to linear type traits and production 1: phenotypic and genetic analysis in pedigree type classified herds. Animal Production (UK). 53: 279-287.
- Campanile G, Di Palo R, De Rosa C, Peretti V, Amante L, Ciotola F and Coletta A (2003). Preliminary results on Mediterranean Italian Buffalo morfometry. Proc.XV ASPA Cong. 337-339.
- CGRFA (Commission on Genetic Resources for Food and Agriculture) 2007. The state of the world's animal genetic resources for food and agriculture final version (Item 3.1(c) of the Draft Provisional Agenda). 11th Regular Session, Rome, (11-15 June).
- Chantala Khana C and Flavey L (1999). Smallholder dairying in the tropic, ILRI (International Livestock Research Institute), Nairobi, Kenya. pp 462.
- Cockrill WR (1974). The Husbadry and Health of

the Domestic Buffalo. Food and Agricultural Organization of the United Nation, Rome (Italy).

- Curik I, Zechner P, Solkner J, Achmann R, Bodo I, Dovc P, Kavar T, Marti E and Brem G (2003). Inbreeding, Microsatellite Heterozygosity, and Morphological Traits in Lipizzan Horses. J. Heredity: 94 (2): 125-132.
- Dario C, Carnicella D, Dario M. and Bufano G (2006). Morphological evolution and heritability estimate for some biometric traits in the Murgese horse breed. Genetic Mol Res. 5 (2): 309-314.
- Dguma G, Mirkena T, Haile A, Iniguez L, Okeyo AM, Tibbo M, Rischkowsky B, Solkner J and Wurzinger M (2010). Participatory approaches to investigate breeding objectives of livestock keepers. Livestock Research for Rural Development 22(4): 1- 13.
- Dia Palo R, Peretti V, De Rosa C, Campanile G and Barbieri V (2001). Computerized image analysis in type trait evaluation of the Mediterranean dairy buffalo (Bubalus bubalis). Proc. XIV ASPA Cong. 225-227.
- Dong Xuan DT, Szalay I, Su VV, Tie HV and Dang Vang N (2006). Animal genetic resources and traditional farming in Vietnam. AGRI. 38: 1-17.
- Doren PE, Baker JF, Long CR and Cartwright (1989). Estimating Parameters of Growth Curves of Bulls. J Anim Sci. 67: 1432-1445.
- Dorji T, Tshering L and Rai DB (2009). Conservation of the last Himalayan cattle breed of Bhutan, South Asia; Pro Poor Livestock Policy Programme; A joint initiative of NDDB and FAO, NDDB house (6th floor) PB 4906, Safdarjang enclave, Newdlhi110029, India. www.sapplpp.org
- FAO (1992). The management of global animal genetic resources. Proceedings of FAO Expert Consultation Rome, Italy, April 1992, (Hoges J. edr).
- FAO (2004). http://appsfao.org/accessed 20 December 2004.
- FAO (2010). LPP, LIFE Network, IUCN-WISP, and FAO (2010). Adding value to livestock diversity-Marketing to promote local breeds. FAO- Animal Production and Health Paper. No. 168. Rome.
- FAO/UNEP (1992). In Situ Conservation of Livestock and Poultry. (Edited: Henson E L), M-22, ISSN 92-5-103143-6; Food and

Agricultural Organization of the United Nation Rome.

- Fernandez G, Baro JA, de la Fuente FL and San Primitivo F (1997). Genetic parameters for linear under traits of dairy ewes. J Dairy Sci. 80: 601-605.
- Gallo I, Carnier P, Cassandro M, Dal Zotto R and Bittante G (2001). Test-Day genetic analysis of condition score and heart girth in Holstein Friesian cows. J Dairy Sci. 84: 2321-2326.
- Gatesy J and Arctander P (2000). Hidden morphological support for the phylogenetic placement of Pseudoryx ngetinheusis with bovine bovids: a combined analysis of gross anatomical evidence and DNA sequences from five genes. Systematic Biology. 49 (3): 515-538.
- Goe MR, Alldredge JR and Light D (2001). Use of heart girth to predict body weight of working oxen in the Ethiopian highlands. Livestock Production Science. 69 (2): 187-195.
- GoP (2006). Livestock Census, 2006. Agricultural Census Organization, Statistic Division, Government of Pakistan, Lahore.
- GoP (2007). Economic Survey of Pakistan, 2006-2007. Finance Division, Government of Pakistan, Islamabad.
- GoP (2009). Economic Survey of Pakistan, 2008-2009 Finance Division, Government of Pakistan, Islamabad
- Hassen F, Bekele E, Ayalew W and dessie T (2007). Genetic variability of five indigenous Ethiopian cattle breeds using RAPD markers. African J Biotech. 6 (19): 2274-2279.
- Islam MR, Saadullah M, Howlidaer AR and Huq MA (1991). Estimation of live weight and dressed carcass weight from different body measurements in goats. Indian J Anim Sci. 61 (40): 460-461.
- Jabbar G (1987). Live weight of Azikheli buffalo at different age groups. J Anim Health Prod. 7: 128-133.
- Janssens S and Vandepitte (2004). Genetic parameters for body measurements and linear type traits in Belgain bleu de mairie, Suffolk and texel sheep. Small Ruminant Research. 54(1): 13-24.
- Jogi RVK and Patil UG (1990). Various body measurements and their correlation with milk yield and fat percentage. Studies in Surti buffaloes. Buffalo Bulletin. 9 (2): 35-38.
- Khan BB, Younas M and Hanjra SH (1982). Breeds

and types of livestock in Pakistan. Department of Livestock Management, University of Agriculture Faisalabad, Pakistan.

- Khan MA (2009). Buffalo the animal of future. 1st, (Edi). Published by Idara Matbuaat-e-Sulemani.
- Khan MA and Niamatullah M (2010). Buffalo versus cattle? Let us close this controversy and concentrate on improving the productivity of buffalo. Proceedings of the 9th World Buffalo Congress Argentina, pp: 1043-1045.
- Khan MS, Ahmad N and Khan MA (2007a). Genetic resources and diversity in dairy buffaloes of Pakistan. Pak Vet J. 27 (4): 201-207.
- Khan MS, Khan MA and Mahmood S (2008b) Genetic resources and diversity in Pakistani goats. International J Agri Biol. 10(2): 227-31.
- Khan MS, Khan MA, Ahmad S and Mahmood S (2007b). Genetic resources and diversity in Pakistani sheep. International J Agri Biol 9 (6): 941-44.
- Khan MS, Rehman Z, Khan MA and Ahmad S (2008a). Genetic resources and diversity in Pakistani cattle. Pak Vet. J. 28 (20): 95-102.
- Kohler-Rollefson I (2001). Intellectual property rights regime necessary for traditional livestock raisers. Indigenous Knowledge Development Monitor. 9 (1): 12-15.
- Kohler-Rollefson I, Rathore HS and Mathias E (2009). Local breeds, livelihood and livestock keepers' rights in South Asia. Trop Anim Health Prod 41: 1061-1070.
- Kolte AY and Sadekar RD (1996). Phenotypic characters of Gauli Strain of Nagpuri (Berari) buffaloes. Indian Vet J. 73: 1094-1096.
- Kumar PP and Raj TM (2007). Gaddi Buffalo: An Indigenous Breed of Far-west Nepal. Ital. J Anim Sci. 6 (Suppl. 2): 1230-1233.
- Lau CH, Drinkwater RD, Yusoff K, Tan SG, Hetzel DJS and Barker JSF (1998). Genetic diversity of Asian water buffalo (Bubalus bubalis): mitochondrial DNA D-loop and cytochrome b sequence variation. Animal genetics
- Lazzaroni C and Bigini D (2005). Influence of rearing system on body measurements in piemontese young bulls. Italian j Anim Sci. 4 (Suppl. 2): 287.
- Loftus T, Machugh DE, Bradley DG, Sharp PM and Cunningham P (1994). Evidence for two

independent domestications of cattle. Proc. National Acad. Sci., USA. 91:2757-2761.

- Luo MF, Wiggans GR and Hubbard SM (1997). Variance component estimation and multitrait genetic evaluation for type traits of dairy goats. Journal of Dairy Science, 80: 594-600.
- Moioli B and Borghese A (2005). Buffalo Breeds and Management Systems: Chapter III: In: Buffalo Production and Research". (Borghese A, edr). FAO Regional Office for Europe, REU Technical Series 67, the state of art of the research, development, products and market of buffalo species in the world.
- Ndumu DB, Baumung R, Wurzinger M, Drucker AG, Okeyo AM, Semambo D and Solkner J (2008). Performance and fitness versus phenotypic appearance in the African Ankole Longhorn cattle: a novel approach to identify selection criteria for indigenous breeds. Livestock Science. 113: 234-242.
- Patil SS, Kale AM and Awati AA (1998). Pandharpuri buffaloes for better persistence. Buffalo Bulletin. 17 (3): 65-66.
- Patro BN, Mishra PK and Rao PK (2008). Chilika buffaloes in Orissa: a unique germplasm. File://F\My-Buff-4.12.8\Morphometry.htm.
- Patro BN, Mishra PK and Rao PK (2008). Chilika buffaloes in Orissa: a unique germplasm. File://F\My-Buff-4.12.8\Morphometry.htm.
- Pretorius SM, Marle-Koster EV and Mostert BE (2004). Description of the Friesian Horse population of South Africa and Namibia. S African J Anim Sci. 34(3): 149- 157.
- Ranjhan SK and Pathak NN (1993). Textbook on buffalo production (Third edition). Vikas Publishing House Pvt LTD, 576, Masjid Road, Jangpura, New Delhi-110014.
- Rege JEO (2001). Defining livestock breeds in the context of community-based management of farm animal genetic resources. In: Proceedings of a Workshop on Communitybased management of Animal Genetic Resources- A Tool for Rural Development and Food Security, Mbabane, Swaziland, (FAO, Rome). 27-35.
- Riva J, Rizzai R, Marelli S and Cavalchinni LG (2004). Body measurements in Bergamasca sheep. Small Ruminant Research. 55: 221-227.
- Roth J and Mayers P (2004). Bubalus bubalis, Animal Diversity Web, viewed 4 July 2005; http://animaldiversity.ummz.umich.edu/site/

accounts/information/bubalus\_buba lis.html

- Scarpa R, Drucker A, Anderson S, Ferraes-Ehuan N, Gomez V, Risoparoin C and Rubio-Leonel O (2003). Valuing animal genetic resources in peasant economies. The case of the Box Keken creole pig in Yuctan. Ecol Economics. 45: 427-443.
- Scherf B, Rischkowsky B and Hoffmann I (2005). Status of animal genetic resources: time for action? International Workshop on" Options and Strategies for the Conservation of Farm Animal genetic Resources" AGROPOLIS' Montpellier, France (7-10 November).
- Schneider MdeP, Durr JW, Cue RI and Monardes HG (2003). Impact of type traits on functional herd life of quebec Holsteins assessed by survival analysis. J Dairy Sci. 86: 4083-4089.
- Sethi RK (2009). Genetic improvement of buffaloes for milk production. Pak J Zool. (Suppl. 9): 559-569.
- Shrikhande GB, Kolte AY and Kolte BR (1996). Phenotypic characters of Nagpuri (Berari) buffaloes. Indian Vet J. 73: 1198-119.
- Singh CV and Barwal RS (2010). Buffalo Breeding Research and Improvement Strategies in India. Proceedings of the 9th World Buffalo Congress Argentina, pp 1024-1031.
- Soysal MI, Tuna YT, Gurcan EK, Ozkan E, Kok S, Castellano N, Cobanoglu O and Barone CMA (2007). Anatolian water buffalo's husbandry in Turkey: Preliminary results on somatic characterization. Ital J Anim Sci. 6 (Suppl. 2): 1302-1307.
- Soysal MI, Tuna YT, Gurcan EK, Ozkan E, Kok S, Castellano N, Cobanoglu O and Barone CMA (2007). Anatolian water buffalo's husbandry in Turkey: Preliminary results on somatic characterization. Italian Journal of Animal Science, 6 (Suppl. 2): 1302-1307.
- Sukla S, Yadav BR and Bhattacharya TK (2006). Characterization of Indian Riverine Buffaloes by Microsatellite Markers. Asian-Australasian Journal of Animal Science. 19 (11): 1556-1560.
- Swaminathan MS (1988). Conservation of animal genetic resources. Proceeding II-World Buffalo Congress Vol. 2 part I: 375.
- Terzano GM, Neglia G, Maschio M, Barile VL, Razzano M, Martiniello P, Cannone I and Borghese A (2007). Effect of intensive or extensive systems on buffalo heifers'

performances: onset of puberty and ovarian size. Ital J Anim Sci. 6 (Suppl. 2): 1273-1276.

- Vij PK and Tantia MS (2005). Status of Nili-Ravi buffaloes in India. AGRI. 37: 75- 81.
- Zarate AV (1996). Breeding strategies for marginal regions in the tropics and subtropics. Anim Res Dev. 4 (44): 99-118.
- Zechner P, Solkner J, Bodo I, Druml T, Baumung R, Achmann R, Marti E, Habe F and Brem G (2002). Analysis of diversity and population structure in the Lipizzan horse breed based on pedigree information. Livestock Prod Sci. 77: 137-146.
- Zechner P, Zohman F, Solkner J, Bodo I, Habe F and Marti E (2001). Morphological description of the Lipizzan horse population. Livestock Prod Sci. 69(2): 163-177.
- Zechner P, Zohman F, Solkner J,Bodo I, Habe F and Marti E (2001). Morphological description of the Lipizzan horse population. Livestock Prod Sci. 69 (2): 163-177.
- Zulu D (2008). Genetic characterization of Zambian Native Cattle Breeds. MSc, thesis (Animal and Poultry Sciences), Faculty of the Virginia Polytechnic Institute and state University, Blacksburg, Virginia.

\*\*\*\*\*